

Ground State Energy for Massive Fields and Renormalisation

M Bordag

Institute for Theoretical Physics, University of Leipzig,

Augustusplatz 10, 04109 Leipzig, Germany

file is mfr2

February 1, 2008

Abstract

We discuss the renormalisation of the ground state energy of massive fields obeying boundary conditions, i.e., of the Casimir effect, and emphasise the role of the mass for its understanding. This is an extended abstract of a talk given at the topical group meeting on Casimir Forces at the Harvard-Smithsonian Center for Astrophysics on March 15-29, 1998.

Renormalisation is a key to the understanding of the structure of quantum field theory. The kind of ultraviolet divergencies occurring divide the perturbative field theoretical models into renormalizable and non or super renormalizable ones. The ultraviolet divergencies occurring during the calculation of ground state energy in different backgrounds (including boundary conditions, i.e., the Casimir effect) carry information on the classical system which one is forced to associate in order to remove resp. interpret the divergencies.

In the present note which is an extended abstract of a talk¹ we discuss the renormalisation using different examples. These are the Casimir effect for massive scalar [1] and spinor [2] fields and the radiative corrections [3] to the electromagnetic Casimir effect with boundary conditions on a sphere. In addition we consider the ground state energy of a scalar field in a spherically symmetric smooth background field [4]. For all questions and references not given in this note we refer to the cited papers.

The necessity to associate some classical system with any ground state energy arises from its very nature. The ground state energy is the amount of energy left in a quantised system when all excitations are gone. To any excited level it

¹given at the topical group meeting on Casimir Forces at the Harvard-Smithsonian Center for Astrophysics on March 15-29, 1998.

adds the same amount of energy (or other observables like charge) and, hence, cannot be observed in any measurements involving a mere change of the quantum numbers (for example in transitions between different levels). The only way to observe ground state energy is to change parameters which are external to the quantised system. In the most prominent example, the Casimir effect between conducting plates, this external parameter is the distance between the plates. Therefore it is inevitable to introduce an appropriate classical system.

The consideration of parallel plates is to some extent misleading. The point is that it is a too simple example, hiding most of the classical structures which one is forced to introduce in the general case. From a very general point of view it is the missing curvature in the case of parallel plates which makes the interesting divergent contributions vanish. What is left is the so called Minkowski space contribution which is independent on the distance between the plates so that one obtains a finite force without any further renormalisation. As a result of this simple behaviour there have been attempts to extend this to the sphere using for example two concentric spheres in order to extract a finite force. Also, efforts have been undertaken to use the remarkable properties of the Zeta functions in order to get infinity free results. As we will argue in the present note, this cannot be successful in the general case.

We consider the following models²:

1. a 'smooth'³ background field. The action is given by

$$S = \frac{1}{2} \int dx \left\{ \Phi(x)(\square - M^2 - \lambda\Phi(x)^2)\Phi(x) + \varphi(x)(\square - m^2 - \lambda'\Phi(x)^2)\varphi(x) \right\}, \quad (1)$$

where $\Phi(x)$ is the classical background field (we choose it static and spherically symmetric) and $\varphi(x)$ is to be quantised. The complete energy of that system is

$$\begin{aligned} \mathcal{E} &\equiv \mathcal{E}_{\text{class}} + \mathcal{E}_{\text{qu}} \\ &= \frac{1}{2} \int d\vec{x} \left((\nabla\Phi(x))^2 + M^2\Phi(x)^2 + \lambda\Phi(x)^4 \right) + \frac{1}{2} \sum_{(n)} \left(\lambda_{(n)}^2 + m^2 \right)^{\frac{1}{2}}, \end{aligned} \quad (2)$$

where the one particle energies are defined by the eigenvalue problem

$$(-\Delta + \lambda'\Phi(\vec{x})^2)\varphi_{(n)}(\vec{x}) = \lambda_{(n)}^2\varphi_{(n)}(\vec{x}). \quad (3)$$

²We use units where $\hbar = c = 1$.

³The word 'smooth' means here the opposite of boundary conditions. In fact, the only example which we consider up to the end is a square well potential which is not smooth in the mathematical sense.

2. a scalar field with boundary conditions on a sphere

$$(\square - m^2)\varphi(x) = 0, \quad \varphi(x)|_{r=R} = 0. \quad (4)$$

3. a spinor field in a bag

$$(i\gamma^\mu \frac{\partial}{\partial x^\mu} + m)\psi(x) = 0, \quad (1 + i\vec{n}\vec{\gamma})\psi(x)|_{r=R} = 0. \quad (5)$$

The last two examples can be subdivided with respect to the region where we consider the quantum field:

- i)* $0 \leq r \leq R$ interior,
- ii)* $R \leq r < \infty$ exterior,
- iii)* $0 \leq r \leq R \cup R \leq r < \infty$ both regions.

The third case, *iii)*, looks like the simple union of *i)* and *ii)*. It is, in fact, with respect to the spectrum of the quantum system. Below we will see how after renormalisation a difference occurs.

The ground state energy is given in all cases by the formula

$$\mathcal{E}_{\text{qu}} = \pm \frac{1}{2} \sum_{(n)} \left(\lambda_{(n)}^2 + m^2 \right)^{\frac{1}{2}},$$

where the sum runs over the spectrum of the corresponding quantum field. The sign in front distinguishes between bosonic and fermionic fields. In *2.i)* we have $(n) = (n, l, m_l)$ and $\lambda_{(n)} = j_{\nu, n}/R$ with the roots of the Bessel functions, $J_\nu(j_{\nu, n}) = 0$ ($\nu \equiv l + \frac{1}{2}$). In *3.i)* the spectrum is determined by the equation

$$J_{\nu+1}(\omega R) + \sqrt{\frac{E-m}{E+m}} J_\nu(\omega R) = 0 \rightarrow \omega = \lambda_{(n)}$$

with $E = \sqrt{m^2 + \omega^2}$. For *ii)* we have to consider some large but finite volume first in order to get a discrete spectrum. By a well known procedure which is explained in detail in [1] one lets that volume tend to infinity throwing away the so called Minkowski space contribution.

In the last two examples the associated classical (geometrical) system has the energy

$$\mathcal{E}_{\text{class}} = pV + \sigma S + FR + k + \frac{h}{R}, \quad (6)$$

where $V = \frac{4}{3}\pi R^3$ is the volume and $S = 4\pi R^2$ is the surface of the sphere. Correspondingly, p is the pressure and σ is the surface tension. The parameters F , k and h do not have a special meaning. This formula is the most general one which can be written down for dimensional reasons. It turns out that this form is required in the cases *i)* and *ii)*, while in *iii)* the first, the second and the third terms can be dropped, see below.

The expression for the ground state energy written so far is divergent and must be regularised. Because of its technical advantages (and its beauty) we use the zeta functional regularisation and write

$$\mathcal{E}_{\text{qu}} = \pm \frac{\mu^{2s}}{2} \sum_{(n)} (\lambda_{(n)}^2 + m^2)^{\frac{1}{2}-s} \quad (7)$$

with sufficiently large $\Re s$ to make the sum converge. In the end one has to perform the analytic continuation to $s = 0$. In Eq. (7) the parameter μ with dimension of a mass was introduced. It is arbitrary and similar to the subtraction point in the renormalisation of perturbative quantum field theory. After renormalisation the ground state energy will become independent of μ .

Eq. (7) by means of

$$\mathcal{E}_{\text{qu}} = \pm \frac{\mu^{2s}}{2} \zeta_{\mathbf{P}}(s - \frac{1}{2}) \quad (8)$$

is in fact the expression of the ground state energy using the zeta function of a corresponding operator $\mathbf{P}$

$$\zeta_{\mathbf{P}}(s) = \sum_{(n)} e_{(n)}^{-s}, \quad (9)$$

where $e_{(n)} = \sqrt{\lambda_{(n)}^2 + m^2}$ are the eigenvalues: $\mathbf{P}\varphi_{(n)} = e_{(n)}\varphi_{(n)}$.

Later on it will be instructive to have another regularisation. We use

$$\mathcal{E}_{\text{qu}} = \pm \frac{1}{2} \sum_{(n)} (\lambda_{(n)}^2 + m^2)^{\frac{1}{2}} e^{-\epsilon(\lambda_{(n)}^2 + m^2)} \quad (10)$$

with $\epsilon \rightarrow 0$ in the end.

The regularised ground state energy (7) can be written in the form (see the cited papers for details, for the dropping of the Minkowski space contribution for instance)

$$\mathcal{E}_{\text{qu}} = -\frac{\cos \pi s}{\pi} \mu^{2s} \sum_{l=0}^{\infty} (l + 1/2) \int_m^{\infty} dk [k^2 - m^2]^{\frac{1}{2}-s} \frac{\partial}{\partial k} \ln f_l(ik). \quad (11)$$

Here, all information is contained in the function $f_l(k)$. In the first model it is the Jost function of the scattering problem corresponding to (3), in the second and third models it is expressed by the corresponding combinations of Bessel and Hankel functions:

- 2. *i*): $f_l(k) \rightarrow J_{\nu}(kR),$
- 2. *ii*): $f_l(k) \rightarrow H_{\nu}^{(1)}(kR),$
- 3. *i*): $f_l(k) \rightarrow J_{\nu+1}(kR) + \sqrt{\frac{E-m}{E+m}} J_{\nu}(kR),$
- 3. *ii*): $f_l(k) \rightarrow H_{\nu+1}^{(1)}(kR) + \sqrt{\frac{E-m}{E+m}} H_{\nu}^{(1)}(kR).$

Thereby the integration runs over the imaginary axis starting from im (cf. Eq. (11)). Representation (11) is equivalent to other ones where the mode density or the scattering phase shift enter. The integration over the imaginary axis has some technical advantages. For instance, there are no oscillations in $f_l(ik)$ and the possible bound states which may occur in a background potential like $\lambda'\Phi(\vec{x})^2$ in (3) are included implicitly.

The general structure of the ultraviolet divergencies can be obtained from the heat kernel expansion. For this reason one represents the zeta function in (9) by an integral

$$\zeta_{\mathbf{P}}(s) = \int_0^\infty \frac{dt}{\Gamma(s)} t^{s-1} K(t), \quad (12)$$

where

$$K(t) = \sum_{(n)} e^{-t(\lambda_{(n)}^2 + m^2)} \quad (13)$$

is the (global) heat kernel. Now the ultraviolet divergencies of the ground state energy are determined from the behaviour of the integrand in (12) at the lower integration limit and, hence, from the asymptotic expansion of the heat kernel for $t \rightarrow 0$:

$$K(t) \sim \frac{e^{-tm^2}}{(4\pi t)^{3/2}} \sum_n a_n t^n \quad n = 0, \frac{1}{2}, 1, \dots \quad (14)$$

This expansion is known for very general manifolds, see the book [5] for example. If the underlying manifold is without boundary, only coefficients with integer numbers enter, otherwise half integer powers of t are present. Sometimes logarithmic contributions occur (not shown in (14)), but we were not confronted with them so far.

Inserting the expansion (14) into Eq. (12) we obtain from the coefficients with $n \leq 2$ (the higher coefficients do not contribute to the ultraviolet divergencies)

$$\begin{aligned} \mathcal{E}^{div} = & -\frac{m^4}{64\pi^2} \left(\frac{1}{s} + \ln \frac{4\mu^2}{m^2} - \frac{1}{2} \right) a_0 - \frac{m^3}{24\pi^{3/2}} a_{1/2} \\ & + \frac{m^2}{32\pi^2} \left(\frac{1}{s} + \ln \frac{4\mu^2}{m^2} - 1 \right) a_1 + \frac{m}{16\pi^{3/2}} a_{3/2} \\ & - \frac{1}{32\pi^2} \left(\frac{1}{s} + \ln \frac{4\mu^2}{m^2} - 2 \right) a_2. \end{aligned} \quad (15)$$

Apparently, the contributions from the coefficients $a_{1/2}$ and $a_{3/2}$ are finite while the other contain a pole in $s = 0$. This is a special feature of the zeta functional

regularisation and not the case in other regularisations. So, for example, in the regularisation given by Eq. (10) the corresponding contribution $\mathcal{E}_{\text{ep}}^{\text{div}}$ to the ground state energy reads

$$\begin{aligned} \mathcal{E}_{\text{ep}}^{\text{div}} = & \frac{3}{64\pi^{5/2}} \left\{ \left(-\frac{4}{3} \frac{1}{\epsilon^2} + 2\frac{m^2}{\epsilon} - \frac{m^4}{2} \ln \epsilon \right) a_0 + \frac{\pi}{2} \left(\frac{1}{\epsilon^{3/2}} - \frac{m^2}{\epsilon^{1/2}} \right) a_{1/2} \right. \\ & \left. + \left(\frac{2}{\epsilon} + m^2 \ln \epsilon \right) a_1 + \frac{\pi}{\epsilon^{1/2}} a_{3/2} - \ln \epsilon a_2 \right\}. \end{aligned} \quad (16)$$

Here the contributions of all coefficients are actually divergent.

The coefficients in the first model are well known:

$$a_0 = \mathcal{V}, \quad a_1 = - \int d\vec{x} \lambda' \Phi(\vec{x})^2, \quad a_2 = \frac{1}{2} \int d\vec{x} \lambda'^2 \Phi(\vec{x})^4. \quad (17)$$

For the second model they read:

$$\begin{aligned} a_0 &= \pm \mathcal{V}, & a_{1/2} &= -2\pi^{3/2} R^2, \\ a_1 &= \pm \frac{8}{3} \pi R, & a_{3/2} &= -\frac{\pi^{3/2}}{6}, & a_2 &= \mp \frac{16}{315} \frac{\pi}{R}, \end{aligned} \quad (18)$$

where the upper sign corresponds to the model 2.i), i.e., to the interior and the lower sign to 2.ii). $\mathcal{V}$ is the volume of the underlying manifold, the whole Minkowski space in the first model, the volume of the interior of the sphere $\frac{4\pi}{3} R^3$ resp. the exterior volume in the second model. In the first model the contribution of a_0 is independent of the background potential and dropped. Similar arguments apply to 2.ii).

The alternation of the signs is valid for general, infinitely thin bounding surfaces, not for the sphere alone. When adding up to get *iii*), the corresponding ultraviolet divergencies cancel between inside and outside. This is just the point, where the third case of the second and third models becomes nontrivial as it has a smaller number of singular contributions than *i*) and *ii*) taken individually.

The renormalisation procedure consists simply in subtracting the divergent contributions from the ground state energy and adding them to the classical contribution:

$$\begin{aligned} \mathcal{E} &= \underbrace{\mathcal{E}_{\text{class}} + \mathcal{E}_{\text{qu}}^{\text{div}}}_{\tilde{\mathcal{E}}_{\text{class}}} + \underbrace{\mathcal{E}_{\text{qu}} - \mathcal{E}_{\text{qu}}^{\text{div}}}_{\mathcal{E}_{\text{qu}}^{\text{ren}}} \\ &\equiv \tilde{\mathcal{E}}_{\text{class}} + \mathcal{E}_{\text{qu}}^{\text{ren}}. \end{aligned} \quad (19)$$

The change from $\mathcal{E}_{\text{class}}$ to $\tilde{\mathcal{E}}_{\text{class}}$ can be interpreted as a renormalisation of the parameters of the classical system. In the first model it reads:

$$M^2 \rightarrow M^2 - \frac{\lambda' m^2}{16\pi^2} \left(\frac{1}{s} + \ln \frac{4\mu}{m} - 1 \right),$$

$$\lambda \rightarrow \lambda - \frac{\lambda' m^2}{64\pi^2} \left(\frac{1}{s} + \ln \frac{4\mu}{m} - 2 \right). \quad (20)$$

The divergence associated with a_0 would lead to a renormalisation of a constant addendum to the classical energy. As said above, we drop such a contribution. We would only like to mention, that in a gravitational background this would be a renormalisation of the cosmological constant. It should be remarked, that the kinetic term in the classical action does not undergo any renormalisation.

In the second model, in the cases *i*) and *ii*), the procedure (19) leads to the following substitutions:

$$\begin{aligned} p &\rightarrow p \mp \frac{m^4}{64\pi^2} \left(\frac{1}{s} - \frac{1}{2} + \ln \left[\frac{4\mu^2}{m^2} \right] \right), & \sigma &\rightarrow \sigma + \frac{m^3}{48\pi}, \\ F &\rightarrow F \pm \frac{m^2}{12\pi} \left(\frac{1}{s} - 1 + \ln \left[\frac{4\mu^2}{m^2} \right] \right), & k &\rightarrow k - \frac{m}{96}, \\ h &\rightarrow h \pm \frac{1}{630\pi} \left(\frac{1}{s} - 2 + \ln \left[\frac{4\mu^2}{m^2} \right] \right). \end{aligned} \quad (21)$$

In contrast, for *iii*) we need only

$$\sigma \rightarrow \sigma + \frac{m^3}{24\pi}, \quad k \rightarrow k - \frac{m}{48}. \quad (22)$$

So we conclude, that in *i*) and *ii*) all five contributions, introduced with (6), are necessary in order to interpret the renormalisation as a redefinition of the parameters of the classical system. In contrast, in the case *iii*), only two are required:

$$\mathcal{E}_{\text{class}}^{iii) = \sigma S + k. \quad (23)$$

Of course, more can be introduced depending of the kind of physical systems one has to consider. But this is the minimal set. Sometimes the physical meaning (and real value) of these parameters is unclear.

The procedure of renormalisation is not unique. Besides different regularisations which can be used and which change the details of formulas like (21) (for instance, when using (10) and (16) instead of (7) and (15)), there is an arbitrariness resulting from the arbitrary parameter μ in (15). Also, after the infinite renormalisation (20) or (21), a finite renormalisation of the same kind is still possible. So one is asked to fix some normalisation condition in order to give the ground state energy a unique meaning. As such we suggest the requirement that the renormalised ground state energy must vanish when making the quantum field infinitely heavy, i.e., for $m \rightarrow \infty$:

$$\mathcal{E}_{\text{qu}}^{\text{ren}} \rightarrow 0 \quad \text{for } m \rightarrow \infty. \quad (24)$$

This condition is physically meaningful. In that limit there should be no quantum fluctuations. On the other hand it is complete, i.e., it fixes $\mathcal{E}_{\text{qu}}^{\text{ren}}$ uniquely

because the last coefficient contributing to divergencies and, hence, the last which can cause a non-uniqueness is a_2 (in general $a_{d/2}$ where d is the dimension of the manifold). All lower coefficients (a_n with $n < \frac{d}{2}$) enter proportional to nonnegative powers of the mass.

The uniqueness of the ground state energy is essential when asking for quantities like the Casimir force⁴. A different situation occurs when considering the back reaction problem. There the dynamics of the background itself is included and one has to look for a minimum of the complete energy $\mathcal{E}$ (2) resp. (19) after renormalisation, varying the background (the field $\Phi(\vec{x})$ or the radius of the sphere, for example). Obviously, in that case a finite renormalisation, i.e., adding zero by subtracting something from $\mathcal{E}_{\text{qu}}^{\text{div}}$ and adding it to $\mathcal{E}_{\text{class}}$, makes no difference.

The goal of fulfilling the normalisation condition (24) is achieved by subtracting the complete contribution resulting from the heat kernel coefficients a_n with $n \leq 2$ as done by (20) resp. (21) and (22) in the first resp. second models. In doing so we obtain $\mathcal{E}_{\text{qu}}^{\text{ren}}$ as defined by (19). Now the regularisation can be removed, i.e., the analytic continuation to $s = 0$ can be performed. This is still a nontrivial task because it cannot be done under the sign of the sum and the integral in expressions like (11). One has to use the uniform asymptotic expansion of $\ln f_l(ik)$ for $l \rightarrow \infty$, $k \rightarrow \infty$ with $\frac{k}{l}$ fixed to the required order (l^{-3} in the considered examples). This results in

$$\mathcal{E}_{\text{qu}}^{\text{ren}} = \mathcal{E}_f + \mathcal{E}_{\text{as}} - \mathcal{E}_{\text{qu}}^{\text{div}}, \quad (25)$$

where

$$\mathcal{E}_f = -\frac{1}{\pi} \sum_{l=0}^{\infty} (l+1/2) \int_m^{\infty} dk [k^2 - m^2]^{\frac{1}{2}} \frac{\partial}{\partial k} (\ln f_l(ik) - \ln f^{\text{as}}) \quad (26)$$

is the 'finite' part. Here, due to the achieved convergence one could put $s = 0$ under the sign of the integral. Also, it is possible to integrate by parts yielding the representation

$$\mathcal{E}_f = \frac{1}{\pi} \sum_{l=0}^{\infty} (l+1/2) \int_0^{\infty} dq (\ln f_l(ik) - \ln f^{\text{as}})_{|_{k=\sqrt{q^2+m^2}}}. \quad (27)$$

In the other ('asymptotic') contribution,

$$\mathcal{E}_{\text{as}} = -\frac{\cos \pi s}{\pi} \mu^{2s} \sum_{l=0}^{\infty} (l+1/2) \int_m^{\infty} dk [k^2 - m^2]^{\frac{1}{2}-s} \frac{\partial}{\partial k} \ln f^{\text{as}}, \quad (28)$$

one has to perform the analytic continuation to $s = 0$ which is quite easy now because the structure of $\ln f^{\text{as}}$ is much simpler than that of $\ln f_l(ik)$. After that

⁴Note that it is only for planar geometries where taking the derivative with respect to a distance removes the divergencies.




Figure 1: The functions $f(x)$ in case *i*), *ii*) and *iii*) for the scalar field

the divergences in $\mathcal{E}^{\text{as}}$ must cancel that of $\mathcal{E}_{\text{qu}}^{\text{div}}$ in (25). In general, for $\ln f^{\text{as}}$ one can take the minimal asymptotic contributions as it was done in the cited papers. But it is possible to include more (non-leading) terms, for instance in order to speed up the convergence in $\mathcal{E}_f$. Once this procedure is carried out (for details see the cited papers) the numerical calculation of the ground state energy can be done.

As an example we consider here the result for the second model, i.e., for the scalar field with Dirichlet boundary conditions on a sphere of radius R . For dimensional reasons the result can be written as

$$\mathcal{E}_{\text{qu}}^{\text{ren}} = \frac{1}{R} f(Rm) = m h(Rm), \quad (29)$$

where $f(x)$ and $h(x)$ are dimensionless functions simply connected by $f(x) = xh(x)$. In fact, the functions f resp. h show the dependence of the energy on the mass m resp. on the radius R . The results obtained in [1] for the function $f(x)$ are shown in figure 1. It is interesting to note the maximums in *i*) and *iii*) for some finite mass. In the function h , i.e., after dividing by R , and when passing to the dependence on the radius, the minimum in case *i*) survives. It is shown in figure 2. In the other two cases, the dependence is simply monotonously decreasing.

The same analysis as for the scalar case had been done in [2] for the spinor field with bag boundary conditions. The result is almost the same, but in detail different (for instance, with respect to the sign). As an example we mention the dependence of the energy on the radius in the case *i*), shown in figure 3. Here, it is interesting to note the minimum.




Figure 2: The function $h(x)$ in case $i)$ for the scalar field




Figure 3: The function $h(x)$ in case $i)$ for the spinor field

The massless case deserves a special consideration. It was not considered in the cited papers [1, 2, 4]. It is known [7] to exhibit the conformal anomaly which is proportional to a_2 so that in case $a_2 \neq 0$ the function $f(Rm)$ in (29) must contain a divergent contribution proportional to $\ln(RM)$ for $m \rightarrow 0$. Therefore, the normalization condition (24) cannot be extended to the massless case. One is left with the fact that the ground state energy cannot be uniquely defined. For example, in the model 2, i) for $m = 0$ it takes in zetafunctional regularisation the form

$$\mathcal{E}_{\text{qu}} = \frac{-a_2}{16\pi R} \left(\frac{1}{2s} + \ln(R\mu) - 1 \right) + \frac{\tilde{h}}{R},$$

where $\tilde{h}$ is some number. The classical contribution must be taken as

$$\mathcal{E}_{\text{class}} = \frac{h}{R}$$

in order to absorb the pol term. But a contribution of the kind $\tilde{h}/R$ resulting from the calculation of the ground state energy has the same dependence on R as $\mathcal{E}_{\text{class}}$ and, hence, it has no predictive power. It suffers also from the arbitrariness resulting from the parameter μ .

The situation is different in case $iii)$. There the divergencies in the $\frac{1}{R}$ -contribution cancel and there is no longer the need to include $\frac{h}{R}$ into the classical energy (cf. (23)). The result

$$\tilde{h} = 0.00282 \tag{30}$$

is unique and has a physical meaning. Therefore, of course, it coincides with earlier calculations.

The just given considerations are of relevance for the electromagnetic Casimir effect on a sphere. The field is massless and for the second photon polarisation which results in Robin boundary conditions the formulas are essentially the same (the heat kernel coefficients a_n take different values, see [6] for example).

Considering an infinitely thin conducting spherical shell, the effect is uniquely defined. If, however, one considers a spherical shell of finite thickness with no field inside we have the cases *i)* and *ii)* with different radii. No cancellation of divergencies between inside and outside occurs. In that case it seems impossible to give a physical meaning to the Casimir energy.

We add just another remark concerning the electromagnetic Casimir effect for a thin spherical shell. In that case the divergent contribution to the energy is zero in the zeta-functional regularisation. Therefor one can obtain the finite result without any renormalisation. In contrast, in different regularisations, two kinds of divergencies are present, for example from (16) we obtain

$$\mathcal{E}_{\text{qu}}^{\text{div}} = \frac{3}{32\pi^{3/2}} \left(\frac{a_{1/2}}{2\epsilon^{3/2}} + \frac{a_{3/2}}{\epsilon^{1/2}} \right). \quad (31)$$

Now, $a_{3/2}$ is independent of the radius R and can be removed by arguments like saying that only the force or the difference in the energy between two conducting spheres of different radii has a physical meaning. The remaining coefficient ($a_{1/2} = -2\pi^{3/2}R^2$ for Dirichlet boundary conditions) turns out to cancel when adding the second photon polarisation which corresponds to Robin boundary conditions. This is the reason why it is possible to obtain a finite result for the electromagnetic Casimir effect on a sphere in other than the zeta-functional regularisations too. This cancelations of divergencies was observed already in the first calculation made by Boyer. Nevertheless it is a rather special feature and does not survive for instance when including radiative corrections. As it was shown in [3], the radiative corrections to the Casimir effect for a thin conducting spherical shell yield a divergent contribution

$$\mathcal{E}_{\text{qu}}^{\text{div}} = -\frac{16}{9\pi}\alpha m_e^3 R^2 - \frac{4}{15\pi}\alpha m_e, \quad (32)$$

where α is the fine structure constant and m_e is the electron mass. While the second term could again be considered as a constant, the first did not cancel between the two polarisations of the photon.

I would like to thank the organisers of the meeting for kind hospitality and G. Barton, K. Kirsten and V. Mostepanenko for stimulating discussions.

References

- [1] M. Bordag, E. Elizalde, K. Kirsten, S. Leseduarte, Phys. Rev. D **56**, 4896 (1997).
- [2] E. Elizalde, M. Bordag, K. Kirsten, J. Phys. A: Math. Gen. **31**, 1743 (1998).
- [3] M. Bordag, J. Lindig, Phys. Rev. D to appear, hep-th/9801129.
- [4] M. Bordag, K. Kirsten, Phys. Rev. D **53**, 5753 (1996).
- [5] P.B. Gilkey, *Invariance Theory, the Heat Equation, and the Atiyah-Singer-Index Theorem* (Publish or Perish, Vilmington, 1984).
- [6] M. Bordag, E. Elizalde, and K. Kirsten, J. Math. Phys. **37**, 895 (1996).
- [7] S.K. Blau, M. Visser and A. Wipf, Nucl. Phys. B, **310**, 163 (1988).